

VARIATION OF TUBER YIELDS AND QUALITY AT SOME JERUSALEM ARTICHOKE GENOTYPES IN PEDOCLIMATIC CONDITIONS FROM CENTER OF MOLDOVA AND THE PLAIN OF OLTENIA, ROMANIA

Simona-Florina Isticioaia¹, Livia Apostol², Gheorghe Matei^{3*}, Oana Mîrzan¹, Paula Pintilie¹, Elena Troțuș¹, Alexandra Leonte¹, Valentin Vlăduț⁴, Oana Cristea⁴, Bogdan Oprea⁵, Bianca Furdui⁶, Cristina Mihaela Marinciu⁷, Marcel Bularda^{6*}

¹Agricultural Research-Development Station Secuieni, 377 Principală str., 617415 Secuieni, Neamț County, Romania

²National Research and Development Institute for Food Bioresources Bucharest, 5 Ancuța Baneasa str., 020323 Bucharest, Sector 1, Romania

³University of Craiova, 13 A.I. Cuza str., 200585 Craiova, Dolj County, Romania

⁴National Research and Development Institute for Machinery and Installations for Agriculture and Food Industry Bucharest, 6 Ion Ionescu de la Brad Blvd., 200396 Bucharest, Sector 1, Romania

⁵National Research and Development Institute for Pedology, Agrochemistry and Environmental Protection Bucharest, 61 Mărăști Blvd., 11464 Bucharest, Sector 1, Romania

⁶“Dunărea de Jos” University of Galați, 47 Domnească str., 800008 Galați, Galați County, Romania

⁷National Agricultural Research and Development Institute Fundulea, 915200 Fundulea, Călărași County, Romania

*Corresponding authors. E-mail: matei.gheorghe@gmail.com; marcelbularda@yahoo.com

ABSTRACT

Existing information in the literature on the variation in tuber yield and their inulin content, depending on the variety and soil and climatic conditions of *Helianthus tuberosus* L., is limited. For this reason, in the pedoclimatic conditions from ARDS Secuieni (Center of Moldova) and from ARDS Caracal (Oltenia Plain) was conducted a study regarding this aspects. This study was conducted between 2018-2020, within an experience with four genotype of Jerusalem artichoke and specifically: Dacic dwarf, Rareș, Olympic and Dăbuleni population. The results obtained showed major differences between the cultivated genotypes and between the two agricultural areas, both in terms of tuber yield and tuber quality. Jerusalem artichoke is a species that has resistance to drought and high temperatures, achieving average yields of 21.3 to ha⁻¹ in Secuieni and 24.0 to ha⁻¹ in Caracal. The Oltenia Plain offered more favorable conditions for the growth and development of Jerusalem artichoke, but the species also successfully succeeded in Central of Moldova. Inulin was present in the tubers in a percentage between 48.35% and 58.38%, which confirms the high functional potential of the species and recommends the species as a “source of fiber”. Also, due to the high mineral content identified in the tubers, the mention of “iron source” and “magnesium source” may be issued.

Keywords: climate, genotype, inulin, production, soil, tuber.

INTRODUCTION

Topinambur or Jerusalem artichoke is a fascinating species with an exceptional history (Kays and Nottingham, 2007).

The species belongs to the *Helianthus* L. genus, *Asteraceae* family, *Asterales* order and has the scientific name of *Helianthus tuberosus* L. (Salaman, 1940).

The second popular name of the species, very often used, the Artichoke of Jerusalem, has nothing to do with the artichoke, this being part of the genus *Cynara* nor with

Jerusalem, appeared due to the description made by Samuel de Champlain, the first European to describe the Jerusalem artichoke from the New World (Massachusetts in 1605), who compared the flavor to that of artichoke in a description of his visit to the homes of the natives with Seigneur De Monts, the leader of the expedition (Bourne, 1906).

Over the last 300 years, interest in this species has varied from area to area. Currently, the latest studies highlight the increased interest in Jerusalem artichoke due to its benefits when included in human and

animal diets and due to its increased potential to be used for biofuel production (Rakhimov et al., 2003).

The Jerusalem artichoke tubers have in their composition 80% water, 15% carbohydrates and 1-2% protein (Fineli, 2004). Data on its composition are relatively scarce compared to other species, however, they reveal significant variations between certain parameters. They contain very little or no starch, are practically fat-free and have a relatively low calorie level (Chaimala et al., 2020). From the small amount of fat in the tubers, unsaturated fatty acids were identified and no saturated fatty acids were determined (Whitney and Rolfes, 1999). Tubers are a good source of dietary fiber, due to the presence of inulin, which is the main storage carbohydrate (Somda et al., 1999). Inulin from tubers varies from 7 to 30% by fresh weight and about 50% by dry weight (Van Loo et al., 1995).

Jerusalem artichoke, native to the north-central part of the USA, is a perennial species that is cultivated as an annual species. It is a temperate zone culture, with a high ecological plasticity, it is cultivated between 40° and 55° north and south latitude (Kays and Nottingham, 2007).

Jerusalem artichoke can be grown on all types of soil, even the poorest in terms of nutrition with minimal cultivation technology. It is considered to be a species with relatively high tolerance to water stress and very high adaptability to unfavorable factors - drought resistance, at extremely high temperatures (Ciuciuc et al., 2019). However, crop productivity is considerably influenced by applied agronomic practices, namely: variety selection, planting date, effective weed control, fertilization, irrigation and harvesting procedures (Slimestad et al., 2010). Most studies suggest planting Jerusalem artichoke in early spring at a depth of 10-15 cm, at a distance between rows between 45-120 cm and at a distance between tubers in a row of

30-60 cm. The optimum soil temperature for planting is between 6 and 7°C (Duke, 1983).

Climate change in Romania forces us to introduce new species in culture, from the perspective, one of them being Jerusalem artichoke. In this species it is necessary to identify and use well-adapted varieties in order to make a faster introduction into production (Khristich et al., 2021) and later in human nutrition. In this sense, in the pedoclimatic conditions from ARDS Secuieni and from ARDS Caracal, in the period 2018-2020, research was conducted on the adaptability of some native varieties/genotypes of Jerusalem artichoke to zonal conditions and on the quality of tubers harvested from experimental fields.

MATERIAL AND METHODS

In order to elucidate the proposed aspects, in the period 2018-2020, a single-factor experiments were organized whose follow-up factor was the genus of Jerusalem artichoke.

The experiments were carried out in the experimental fields of the Secuieni and Caracal Agricultural Research-Development Stations, according to the method of the Latin rectangle, in three repetitions.

The soil

At ARDS Caracal, the location of the experiments was made on an argic chernozem, in which the particle size composition showed that the soil falls into the group of fine textures, the soil being clay - medium clay, and at ARDS Secuieni on a typical phaeozem (chernozem) cambic soil type, with medium texture. In Caracal the soil is characterized as weakly alkaline, well to very well supplied with potassium, weak to medium fertile and poorly supplied with nitrogen, and in Secuieni, as weakly acidic, well to very well supplied with mobile phosphorus and potassium, poorly supplied with nitrogen and poorly fertile (Table 1).

SIMONA-FLORINA ISTICIOAIA ET AL.: VARIATION OF TUBER YIELDS AND QUALITY
AT SOME JERUSALEM ARTICHOKE GENOTYPES IN PEDOCLIMATIC CONDITIONS
FROM CENTER OF MOLDOVA AND THE PLAIN OF OLTENIA, ROMANIA

Table 1. Characterization of the type of soil on which the experiments were located

Soil characteristics	ARDS Caracal		ARDS Secuieni	
	Argic chernozem		Phaeozem (chernozem) cambic	
	Value	Interpretation	Value	Interpretation
Humus content	3.2%	weak to medium fertile	2.3%	weak fertile
Total nitrogen content	0.130%	poorly supplied with nitrogen	0.134%	poorly supplied with nitrogen
Affordable phosphorus content	44 ppm	medium to well supplied with phosphorus	74 ppm	well to very well supplied with mobile phosphorus
Affordable potassium content	234 mg/kg	good to very well supplied with potassium	221 mg/kg	good to very well supplied with potassium
pH	7.6	weak alkaline	6.1	weak acid

From a climatic point of view, the period 2018-2020 was characterized as warm in both study areas, dry in the Secuieni area and normal to rainy in the Caracal area.

Comparing the two agricultural areas, the Oltenia Plain was warmer and richer in precipitation compared to the Center of Moldova (Figure 1).




Figure 1. Climatic conditions registered in the experimental areas, 2018-2020

The biological material used in the experiment was provided by Vegetable Research and Development Station Buzău and Research and Development Station for Agricultural Plants on Sands Dăbuleni, represented by the varieties Rareș, Olympic, Dacic dwarf and the Dăbuleni local population.

The Jerusalem artichoke was planted in the field in the spring, the third decade of March - the first decade of April, at a distance of 70 cm between rows and 50 cm between tubers/row. The sowing depth was 10 cm, and the previous plant was soybeans.

The tubers were harvested in late October - early November.

In the, within the, The physico-chemical and nutritional characterization of these varieties was performed at laboratory of the National Research-Development Institute for Food Bioresources Bucharest. This unit using modern methods of analysis, apparatus and equipment for analysis, measurement and processing specific to raw materials and products (<https://erris.gov.ro/Food-Chemistry-Laboratory>; <https://erris.gov.ro/Microbiology-ELISA-Laboratory>; <https://erris.gov.ro/Food-Packaging-Laboratory>).

RESULTS AND DISCUSSION

Following the research carried out in the three agricultural years, it was observed that both the genotype experienced and the pedoclimatic conditions had influences on both the yield and the quality of the Jerusalem artichoke tubers.

Compared to the average experience (control), the Rareș and Olympic varieties achieved statistically assured yield increases in each of the three years of experimentation and in each location. These were interpreted as being very significant for the Olympic variety every year in both locations and for the Rareș variety in the first and second year of vegetation in Secuieni and in the first year in Caracal. The Dacic dwarf varieties and the population of Dăbuleni achieved significant negative yield differences in both locations (Table 2).

Thus, from Table 2 it can be seen that the best conditions for the growth and development of Jerusalem artichoke were those of the third year of experimentation in

Caracal (2020), when an average tuber yield of 36.6 to ha⁻¹ was obtained. In this pedoclimatic area, the second year of experimentation was extremely unfavorable to the crop, obtaining an average production of 10.3 to ha⁻¹.

In Secuieni, the average yield obtained in the first two years of experimentation were similar, these having values of 24.7 to ha⁻¹ (2018) and 24.1 to ha⁻¹ (2019), respectively. The third year of experimentation offered extremely unfavorable conditions for the growth and development of Jerusalem artichoke, the average yield being only 15.0 to ha⁻¹ (Table 2).

Regardless of the pedoclimatic conditions, the Olympic variety stood out as superior to the other varieties, being productively ranked first in each year of experimentation in both locations, with an average yield of 37.6 to ha⁻¹, respectively 32.0 to ha⁻¹ in year I, of 36.6 to ha⁻¹, respectively 14.6 to ha⁻¹ in second year and of 20.6 to ha⁻¹, respectively 47.1 to ha⁻¹ in the third year. This was followed by yield close to the Rareș variety (Table 2).

Table 2. The tuber yield for studied varieties of Jerusalem artichoke (*Helianthus tuberosus* L.) in the period 2018-2020

Variety	Tuber yield (to ha ⁻¹)					
	Year I		Year II		Year III	
	Secuieni	Caracal	Secuieni	Caracal	Secuieni	Caracal
Rareș	32.7 ^{***}	31.9 ^{***}	31.9 ^{***}	11.5 [*]	18.4 ^{**}	41.7 ^{**}
Olympic	37.6 ^{***}	32.0 ^{***}	36.6 ^{***}	14.6 ^{***}	20.6 ^{***}	47.1 ^{***}
Dacic dwarf	17.6 ⁰⁰⁰	27.1 [*]	18.0 ⁰⁰⁰	9.4	9.8 ⁰⁰⁰	27.1 ⁰⁰⁰
Dăbuleni Pop	10.8 ⁰⁰⁰	8.9 ⁰⁰⁰	9.8 ⁰⁰⁰	5.8 ⁰⁰⁰	11.0 ⁰⁰⁰	30.6 ⁰⁰⁰
Average	24.7 ^{Mt}	25.0 ^{Mt}	24.1 ^{Mt}	10.3 ^{Mt}	15.0 ^{Mt}	36.6 ^{Mt}
LSD 5% (to ha ⁻¹)	1.7	1.6	2.3	1.1	1.4	1.6
LSD 1% (to ha ⁻¹)	2.4	2.3	3.1	1.8	2.1	3.4
LSD 0.1% (to ha ⁻¹)	3.7	3.4	5.8	3.2	3.8	5.8

On average over the three years of experimentation, the tuber yield varied from one variety to another and from one area to another, being between 10.5 to ha⁻¹ and 31.6 to ha⁻¹ in the conditions from Secuieni and between 15.1 to ha⁻¹ and 31.2 to ha⁻¹ in the conditions of Caracal. The Olympic variety stood out with superior yields in both experimental areas, these being 31.6 to ha⁻¹ and 31.2 to ha⁻¹, respectively.

Compared to the average experience, the control of the experience, statistically assured

yield increases were obtained for the Olympic and Rareș varieties, these being interpreted as being very significant in Secuieni and distinctly and very significant in Caracal. The other genotypes experienced, namely the Dăbuleni population and Dacic dwarf, achieved very significant negative yield differences in Secuieni and distinct negative and very significant in Caracal (Table 3).

The average yield/experience was 21.3 to ha⁻¹ in the conditions from Secuieni and 24.0 to ha⁻¹ in the conditions from Caracal. This highlights

SIMONA-FLORINA ISTICIOAIA ET AL.: VARIATION OF TUBER YIELDS AND QUALITY
AT SOME JERUSALEM ARTICHOKE GENOTYPES IN PEDOCLIMATIC CONDITIONS
FROM CENTER OF MOLDOVA AND THE PLAIN OF OLTENIA, ROMANIA

a greater favorability of the pedoclimatic conditions in the Caracal area for the cultivation of Jerusalem artichoke, but nevertheless the

species achieved good productions even in the conditions from Secuieni (Table 3).

Table 3. The average of tuber yield for studied varieties of Jerusalem artichoke (*Helianthus tuberosus* L.) in the period 2018-2020

Variety	Yield tubers							
	Secuieni				Caracal			
	to ha ⁻¹	%	Diff. from the control	Semn.	to ha ⁻¹	%	Diff. from the control	Semn.
Rareș	27.7	130	6.4	***	28.4	118	4.4	**
Olimpic	31.6	148	10.3	***	31.2	130	7.2	***
Dacic dwarf	15.1	71	-6.2	ooo	21.2	88	-2.8	oo
Dăbuleni Pop.	10.5	49	-10.8	ooo	15.1	63	-8.9	ooo
Media (control)	21.3	100	Mt.		24.0	100	Mt.	
LSD (to ha ⁻¹)	5% = 2.3; 1% = 3.5; 0.1% = 5.5				5% = 1.7; 1% = 2.8; 0.1% = 4.9			

The results obtained from the physico-chemical and nutritional analysis of the tubers showed that Jerusalem artichoke has in its composition substances with functional potential, namely inulin and crude fibers. Of the four genotypes analyzed, the Olympic variety achieved the highest inulin content in both areas, of 55.41 % and 56.15 %, respectively.

Given the high percentages of inulin (prebiotic) identified in Jerusalem artichoke tubers, higher than 50%, the species is considered a „fiber source” [Regulation (EC) No 1924/2006]. The experimental analysis of the chemical composition of Jerusalem artichoke tubers shows that the total constituent substance is represented by the inulin content

(48.35% and 56.15%) (Table 4). This high inulin content confirms that Jerusalem artichoke tubers of studied varieties have great functional potential. Numerous studies have shown that inulin is a valuable substance for human health, having a beneficial action from the moment it enters the human body through the stomach and up to the gastrointestinal tract (Rubel et al., 2014; Ramnani et al., 2010).

The highest protein content, of 19.58%, was identified in the tubers of the Dacian variety (19.58%), under conditions of Caracal location. The high ash content of the Jerusalem artichoke samples indicates that they have a high mineral content (Table 4).

Table 4. The physico-chemical composition of tubers from studied Jerusalem artichoke varieties

Compound	Mineral content							
	Rareș		Olimpic		Dacic dwarf		Dăbuleni Pop.	
	Secuieni	Caracal	Secuieni	Caracal	Secuieni	Caracal	Secuieni	Caracal
Moisture %	7.03	8.65	7.29	9.18	7.53	8.70	6.26	8.78
Ash %	3.73	3.52	3.95	3.13	3.74	3.84	3.35	3.87
Lipids %	0.39	0.39	0.35	0.31	0.28	0.26	0.31	0.24
Proteins %	12.15	12.58	11.52	11.84	11.66	19.58	12.3	13.77
Crude fiber %	3.08	3.36	3.12	3.28	3.56	3.67	3.00	3.00
Inulin %	55.04	54.67	55.41	56.15	55.04	48.35	54.67	55.04

The minerals determined in the Jerusalem artichoke samples collected from the two locations had values of high contents, as follows (mg/100 g s.u.): phosphorus - 260-360; potassium - 2666-3277; calcium - 13.1-21.3; magnesium - 230-240; iron - 15.1-22.3; zinc -

1.74-2.44; copper - 0.44-1.05; manganese - 0.33-0.84 (Table 5).

The iron content identified in the tubers of the population of Dăbuleni exceeds the recommended daily dose, but the one identified in the rest of the varieties is close

to the recommended daily dose. Also, the magnesium and potassium content identified in the tubers exceeds half of the recommended daily dose. The copper content of the Dacian variety grown at ARDS Caracal exceeds half of the recommended daily dose (Table 5). Following the results obtained, Jerusalem

artichoke should be referred to as „source of iron” and „source of potassium” (Regulation EC no. 1924/2006; Directive 90/496/EEC).

The high mineral content, especially potassium, iron and magnesium, is confirmed by other research in the field (Bach et al., 2013; Somda et al., 1999; Terzic et al., 2012).

Table 5. The mineral composition of tubers from Jerusalem artichoke varieties

The mineral	Mineral content							
	Rareş		Olimpic		Dacic dwarf		Dăbuleni Pop.	
	Secuieni	Caracal	Secuieni	Caracal	Secuieni	Caracal	Secuieni	Caracal
P mg/100g su	260	360	270	290	310	360	260	310
K mg/100g su	3020	2988	2698	2777	3000	2666	3177	3277
Ca mg/100g su	18.9	13.9	14.2	13.1	21.3	17.7	19.7	16.8
Mg mg/100g su	230	240	230	220	230	240	230	230
Fe mg/100g su	15.7	17.3	16.4	15.8	19.3	17.2	22.3	15.1
Zn mg/100g su	1.74	2.30	1.85	1.93	1.98	2.44	1.98	2.27
Cu mg/100g su	0.44	0.66	0.51	0.65	0.84	1.05	0.73	0.88
Mn mg/100g su	0.33	0.70	0.44	0.53	0.49	0.84	0.58	0.62

CONCLUSIONS

The average Jerusalem artichoke yields obtained in Secuieni (21.3 to ha⁻¹) and Caracal (24.0 to ha⁻¹), two localities in Romania with drought and strong heat conditions, demonstrate the drought and heat resistance of the studied varieties. Which suggests that this species may be an alternative in the conditions of climate change in Romania.

Although, the Oltenia Plain offers more favorable conditions for the growth and development of Jerusalem artichoke, the species also succeeds successfully in Central of Moldova.

Inulin was present in the tubers in a percentage between 48.35% and 58.38%, which confirms the high functional potential of the species and recommends the species as a „source of fiber”.

Also, due to the high mineral content identified in the tubers, the mention of „iron source” and „magnesium source” may be issued.

ACKNOWLEDGEMENTS

This work was supported by a grant of the Romanian Ministry of Research and Innovation CCDI - UEFISCDI, „Complex system of integral capitalization of agricultural species

with energy and food potential”, project number PN-III-P1-1.2-PCCDI-2017-0566, Contract no. 9PCCDI/2018, within PNCDI III.

To carry out this study, all authors made equal efforts, as a result all authors have equal rights.

REFERENCES

- Bach, V., Jensen, S., Kidmose, U., Sorensenl, J., Edelenbos, M., 2013. *The effect of culinary preparation on carbohydrate composition. Texture and sensory quality of Jerusalem artichoke tubers (Helianthus tuberosus L.)*. LWT - Food Science and Technology, 54: 165-170.
- Bourne, A.T., 1906. *The voyages and explorations of Samuel de Champlain (1604-1616)*. A.S. Barnes Co., New York.
- Chaimala, A., Jogloy, S., Vorasoot, N., Toomsan, B., Jongrunklang, N., Kesmala, T., Holbrook, C.C., Kvien, C.K., 2020. *Responses of total biomass, shoot dry weight, yield and yield components of jerusalem artichoke (Helianthus tuberosus L.) varieties under different terminal drought duration*. Agriculture 2020, 10: 198. <https://doi.org/10.3390/agriculture10060198>
- Ciuciuc, E., Drăghici, I., Drăghici, R., Croitoru, M., Băjenaru, M.F., 2019. *The behavior of varieties of jerusalem artichoke on the sandy soils from south of Oltenia*. Annals of the University of Craiova - Agriculture, Montanology, Cadastre Series, Vol. XLIX, 52: 57.
- Duke, J., 1983. *Handbook of Energy Crops*. http://www.hort.purdue.edu/newcrop/duke_energy/dukeindex.html.

SIMONA-FLORINA ISTICIOAIA ET AL.: VARIATION OF TUBER YIELDS AND QUALITY
AT SOME JERUSALEM ARTICHOKE GENOTYPES IN PEDOCLIMATIC CONDITIONS
FROM CENTER OF MOLDOVA AND THE PLAIN OF OLTENIA, ROMANIA

- Kays, S.J., and Nottingham, S.F., 2007. *Biology and chemistry of Jerusalem artichoke: Helianthus tuberosus L.* CRC Press, Boca Raton, FL.
- Khrstich, V.V., Frizen, Yu V., Gaivays, A.A., 2021. *Research results of Jerusalem artichoke varieties and hybrids in the forest-steppe of the Omsk region.* IOP Conf. Ser.: Earth Environ. Sci., 624: 012071.
- Rakhimov, D.A., Arifkhodzhaev, A.O., Mezhlumyan, L.G., Yuldashev, O.M., Rozikova, U.A., Aikhodzhaeva, N., Vakil, M.M., 2003. *Carbohydrate and proteins from Helianthus tuberosus.* Chemistry of Natural Compounds, 39: 312-313.
- Ramnani, P., Gaudier, E., Bingham, M., van Bruggen, P., Tuohy, K.M., Gibson, G., 2010. *Prebiotic effect of fruit and vegetable shots containing Jerusalem artichoke inulin: a human intervention study.* British Journal of Nutrition, 104: 233-240.
- Rubel, A., Pérez, E., Genovese, D., Manrique, G., 2014. *In vitro prebiotic activity of inulin-rich carbohydrates extracted from Jerusalem artichoke (Helianthus tuberosus L.) tubers at different storage times by Lactobacillus paracasei.* Food Research International, 62: 59-65.
- Salaman, R.N., 1940. *Why "Jerusalem" artichoke?* J. Royal Hort. Soc., LXVI: 338-348, 376-383.
- Slimestad, R., Seljaasen, R., Meijer, K., Skar, S.L., 2010. *Norwegian-grown Jerusalem artichoke (Helianthus tuberosus L.): morphology and content of sugars and fructo-oligosaccharides in stems and tubers.* J. Sci. Food Agric. 90: 956-964.
- Somda, Z.C., McLaurin, W.J., Kays, S.J., 1999. *Jerusalem artichoke growth, development, and field storage. II. Carbon and nutrient element allocation and redistribution.* J. Plant Nutr., 22: 1315-1334.
- Terzic, S., Atlagic, J., Maksimovic, I., Zeremski, T., Zoric, M., Miklic, V., 2012. *Genetic variability for concentrations of essential elements in tubers and leaves of Jerusalem artichoke (Helianthus tuberosus L.).* Scientia Horticulturae, 136: 135-144.
- Van Loo, J.P., Coussement, P., De Leenheer, L., Hoebregs, H., Smits, G., 1995. *On the presence of inulin and oligofructose as natural ingredients in the Western diet.* Crit. Rev. Food Sci. Nutr., 35: 525-552.
- Whitney, E.N., and Rolfes, S.R., 1999. *Understanding Nutrition.* 8th ed., West/Wadsworth, Belmont, CA.
- ***Council Directive 90/496/EEC of 24 September 1990 on nutrition labelling for foodstuffs. <http://data.europa.eu/eli/dir/1990/496/oj>
- ***Fineli, Food Composition Database, National Public Health Institute of Finland, Helsinki, 2004.
- ***Regulation (EC) No 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods. <http://data.europa.eu/eli/reg/2006/1924/oj>.
- ***<https://erris.gov.ro/Food-Chemistry-Laboratory>
- ***<https://erris.gov.ro/Microbiology-ELISA-Laboratory>
- ***<https://erris.gov.ro/Food-Packaging-Laboratory>